



## Research Article



# Diversity of Butterflies (Insecta: Lepidoptera) in Kasur district, Punjab, Pakistan

Muhammad Amir Sial

Department of Wildlife and Ecology, University of Veterinary and Animal Science, Lahore, Pakistan

\*Corresponding author e-mail: [mramirsyal@gmail.com](mailto:mramirsyal@gmail.com)

(Received: 18/07/2023; Revised: 10/09/2023; Accepted: 05/11/2023; Published: 20/12/2023)

### ABSTRACT

The purpose of the study was to investigate the butterfly diversity in several areas of Pakistan's Punjab Province's Kasur district. This Study was conducted in April to May, 2023 at different localities like Changa Manga Forest, Rana Resort Forest and Pattoki Nurseries of Kasur district. During this survey, 106 butterfly specimens were gathered manually and with the aid of an insect net. Five distinct species from four different genera and two separate families (Nymphalidae Pieridae) were found in these specimens. This is the first study to explore the butterfly diversity in district Kasur.

**Keywords:** Butterfly diversity, Kasur.

### INTRODUCTION

Butterflies belong to order Lepidoptera which is the second largest order of class Insecta. Moths and butterflies serve as flower pollinators and are beneficial to both humans and other animals because they help in the growth of fruits, production and seeds [1]. Butterfly larvae and other insects are the primary plant herbivores, eating the leaves of trees, shrubs, and other plants to obtain energy and build a food chain web [2]. Adult butterflies consume flower nectar, which gives them energy. Pollen, on the other hand, gives butterflies access to vitamins, proteins, and lipids. Caterpillars are the term for a butterfly or moth's larva [3]. Caterpillars are effective pests because they seriously harm a variety of plants by defoliating them [4]. Around 28,000 species of butterflies have been identified globally, with 80% of species found in tropical regions [5]. In Pakistan, there are about 5000 species of insects comprising about 400 species of moths and butterflies [6]. T.J Robert worked on the taxonomy of butterflies from different localities of Pakistan [7]. In the past research was carried out to find the diversity of butterflies at different regions of Punjab like Haripur, Azad Kashmir and Gujrat but this district was untouched regarding butterfly diversity. This is the first effort to explore the butterfly diversity in this area.

### MATERIALS AND METHODS

#### Study Site

The Study Conducted on three different sites which are Changa Manga Forest, Pattoki Nurseries and Rana Resort Forest, at district Kasur, Punjab province, Pakistan. Kasur is a district of Lahore division which is situated on the border of India and Pakistan. It consists

of four Tehsils which are Pattoki, Kot Radaha Kishan, Chnian, Kasur. The total land area of district Kasur is 3,995 km<sup>2</sup>.

#### Changa Manga Forest:

Changa Manga Park (31° 5' 0" N, 73° 58' 0" E) being maintained by the Punjab Forest Department, Kasur Forest Division is one of the very few parks for out-door recreation in Pakistan. British forestry experts planted the first trees in the Changa Manga forest in 1866. It is located 80 kilometres to the south-west of Lahore. It covers an area of about 5,065 hectares. It contains many parks of flowers therefore it is a good habitat for butterflies.

#### Rana Resort Forest:

Opened in 2007, Rana Resort and Safari Park is located next to the Bhai Pheru Head Balloki Road in Nankana Sahib, Punjab. Its coordinates are 31° 13' 48.9" N and 73° 51' 31.57" E. This resort was built with the intention of entertaining Lahore residents and safeguarding the rapidly declining rare species of wildlife. There are many parks therefore it is a good habitat for butterflies. The lavish resort is surrounded by a wildlife sanctuary comprising 14 acres of lush green flower-laden lawns and 200 acres of sprawling bamboo forest.

#### Pattoki Nurseries:

In the outskirts of Lahore, Pakistan, at 31° 1' 0" North and 73° 51' 0" East, is Pattoki, a typical Punjabi rural market town famed for its flower cultivation. One of the largest collections of nurseries for flowers, fruits, and ornamental plants in the nation may be found in the town. This makes it a suitable home for butterflies.

### Collection and Preservation

Beginning in April and continuing through the end of May 2023, the locations were visited every week. The butterflies were gathered during the day, from 9 am to 4 pm, from each location. The specimens were collected by using the insect net and hand picking method. The specimens were collected by usual methods of collecting the butterflies, that is, chasing the butterflies with a net in hand. Soon after collecting the butterflies were killed by using chloroform and stretched on a stretching board, then transferred to the insect boxes. The collected specimens were identified up to the species level with help of standard available literature of already identified species, thesis, research articles [8, 9, 10] and Facebook groups like butterflies of Pakistan. Identified specimens were labelled with their scientific names and location of collection along with date properly.




Figure A. Map of Kasur District (highlighted in red) within Punjab in fig: A.

Figure B. Map of District Kasur depicting the sampling sites, which are highlighted in black colour in fig: B.

### RESULTS AND DISCUSSION

The present studies were conducted on the identification and distribution of butterflies (Class, Insecta; Order, Lepidoptera) of Kasur district. Total of 106 specimens were collected from 3 different localities. The identified specimens of butterflies belonged to 5 species, 5 genera and 2 families and the total number of collected specimens are shown in table 1. Family Nymphalidae comprises the largest number of Species and most prevalent family of 60 (56.6%) followed by Pieridae 46 (43.6%) (Table 2).

Order : Lepidoptera

**Family : Nymphalidae**

Sub Family : Danainae

Genus : *Danaus*

*Danaus chrysippus* (Figure 1)

*Danaus chrysippus* (Linnaeus 1758) also known as the plain tiger, is a medium-sized (roughly the size of an Oreo cookie). Its wings are orange with black margins that are speckled with white. From a distance, its orange-black-white colour scheme is distinctive, earning the tiger its name. The plain tiger is typically observed in open spaces, such as gardens and parks, close to their supporting plants. The entire continent of Africa, Mediterranean Europe, the Middle East, much of Asia, and even Australia have all reported seeing it.

Subfamily : Heliconiinae

Genus : *Phalanta*

*Phalanta phalantha* (Figure 2)

*Phalanta Phalantha* (Drury, 1773) is also known as the common leopard. Its upper surface's background coloured is consistently orange, with varying dark brown spots and streaks spread across the wings. The forewing apex and outer margins are not dark brown. The underside's colour and design can differ. The hue of the background takes on a variety of pale orange hues, with a purplish undertone visible in some newly emerging individuals. On the hindwing of certain individuals, there is additionally a middle band with scattered white spots. Its eyes have a grey colour. It is widespread over Africa and South Asia.

Subfamily : Nymphalinae

Genus : *Junonia*

*Junonia almana* (Figure 3)

*Junonia almana* (Linnaeus 1758) is also known as Peacock pansy. The hindwings of this butterfly have pronounced eyespots and an upperside that is bright orange. The eyespots are still quite noticeable despite the undersides being a little paler. Above, the wings are rich yellowish brown with darker brown edges at both termen and costal margins. It is distributed in India, Sri Lanka, Southeast Asia and further east in China and Japan.



Figure 1.



Figure 2.



Figure 3.



Figure 4.



Figure 5.




Figure 6. Species wise distribution at district Kasur.

**Family : Pieridae**

Subfamily : Coliadinae

Genus : *Eumera*

*Eumera hecabe* (Figure 4)

*Eumera hecabe* (Linnaeus 1758) is also known as common grass yellow. The butterfly's forewing has an outer edge that is broad, black, and irregular, while the hindwing has one that is thin and generally regular. Both its forewing and hindwing have an outer edge that is not entirely wavy or totally rounded. There are several black markings on its underside. On the ends of each cell, there is a mark that is always black with a yellow centre. When this happens, they are smaller than the Three Spot's chocolate-coloured area at the tip of the forewing. A little species of pierid butterfly called *Eumera hecabe*, sometimes known as the common grass yellow, is found in South America, Australia, and Asia.

Subfamily : Pierinae

Genus : *Pieris*

*Pieris brassicae* (Figure 5)

*Pieris brassicae* (Linnaeus 1758) is also known as large white. This butterfly has brilliant white wings and black tips to the forewings, extending down the wing edge. Females have two spots on the forewings, which is not present in males. Two spots are present on the creamy white undersides. The forewings are white with one or two small round black spots and a sizable anterior black patch; the hindwings have a single anterior black mark. The head is blackish in coloured, these butterflies have hairy larvae with green bodies that are dotted with black spots and have yellow lateral stripes.

In district Kasur, a total 106 butterfly specimens were gathered from 3 distinct locations. The identification of these collected specimens show that they belong to five different butterfly species from five (05) genera, two (02) families in table 1. Nymphalidae is the family with the greatest percentage of species (56.6%), followed by Pieridae (43.6%) in table 2. In a previous study, the Haroon gathered 232 butterflies which were 13 species and belonging to 3 families to explore the diversity of butterflies in the union council of Koaz Bahram Dheri, District Charsadda, Khyber Pakhtunkhwa. Most species (49%) belong to the family Nymphalidae, followed by Pieridae (37%) and Papilionidae (14%) [11]. The present study matches up with the above mentioned study because both have the highest percentage of Nymphalidae family. Rahman had explored the diversity at Murree; a total 30 species from 9 families were recorded from Murree. Family Nymphalidae has the highest number of species[12]. A study conducted by Parveen and Haroon during this study, 506 specimens in all, belonging to 3 families, 18 genera, and 23 species, were gathered from Tehsil Tangi in Khyber Pakhtunkhwa. The analysis revealed that the family Nymphalidae has the greatest number of members in the current checklist [13]. Between September and December 2008, 21 species from 3 different families were found in Kohat, Pakistan. The reported families Nymphalidae comprised 33% of the butterfly

biodiversity in Kohat, Papilionidae 10%, and Pieridae 57% and in this study family Pieridae has the highest number of species, this study conducted by Parveen [14]. Rhe-philipe reported 5 families from Lahore, including Papilionidae (3 species), Pieridae (17 species), Lycaenidae (11 species), Nymphalidae (16 species) and Hesperioidea (7 species) and family Pieridae has highest number of species [15]. In the Gujrat district, Iqbal conducted research in various areas. From various locations, 232 adult butterfly specimens were gathered for the study. The identification of twelve species of butterflies from eight genera, four subfamilies, and three families was done using these specimens. A total of 78.44%, 8.18%, and 13.36% of the species in this study region belong to the Pieridae, Nymphalidae, and Danaidae families [16].

The current research represents the first effort to investigate the diversity of butterflies in Kasur district, Punjab, Pakistan. This is a completely new contribution to the already existing literature of the world.

**Table 1.** The number of butterfly species collected from three localities of Kasur district during the study period.

Common Names	Scientific Names	Changa Manga	Pattoki	Rana Resort	Total
Plain tiger	<i>Danaus chrysippus</i>	13	6	9	28
Common leopard	<i>Phalanta phalantha</i>	9	7	6	22
Peacock Pansy	<i>Junonia almana</i>	6	4	0	10
Large White	<i>Pieris brassicae</i>	10	8	8	26
Common Grass Yellow	<i>Eurema hecabe</i>	7	9	4	20
Total		45	34	27	106

**Table 2.** Total number and Percentage of two families collected from the three localities of Kasur district during study period.

Families	Changa Manga	Pattoki	Rana Resort	Total
Nymphalidae	27	17	15	60 (56.6%)
Pieridae	17	17	12	46 (43.3%)

## CONCLUSIONS

From the current study it was concluded that the family Nymphalidae present in higher numbers in district Kasur, Punjab, Pakistan.

## ACKNOWLEDGEMENTS

The author is grateful to Akram Awan, Admin of social media group, Facebook, Butterflies of Pakistan, for identification of butterflies and also gratitude to Officials, Department of Wildlife and Ecology, University of veterinary and animal science, Lahore, Punjab, Pakistan for providing facilities throughout the present study.

## CONFLICT OF INTEREST

The author here declares that there is no conflict of interest in the publication of this article.

## REFERENCES

1. Maheshwari JK 2003. Endangered pollinators. *Environmental News Archives*. **9**(1).
2. Dayananda GY 2014. Diversity of butterfly fauna in and around Gudavi bird sanctuary, Sorab, Karnataka. *Journal of Entomology and Zoology Studies*. **2**(5):376-80.
3. Borror DJ, Delong DM, Triplehor CA. 1975. Order Lepidoptera. An introduction to the Study of Insects, New York. 462-534.
4. Kumar S, Khamashon L, Pandey P, Chaudhary R, Nath P, Awasthi S, Joshi PC. 2013. Community Composition and Species Diversity of Butterfly Fauna within Gurukula Kangri Vishwavidyalaya Campus. *Journal of Entomology and Zoology Studies*. **1**(6):66-69.
5. Khan MI, Ullah H, Suleman S, Khan MA, Muhammad N, Zada S, Hussain S. 2015. A review on diversity of butterfly fauna in Pakistan. *World Journal of Zoology*. **10**(4):313-317.
6. Khan MR, Rafi MA, Munir M, Hussain S, Baig MW, Khan MW. 2007. Biodiversity of Butterflies from Districts Kotli, Mirpur and Bhimber, Azad Kashmir. *Pakistan Journal of Zoology*. **39**(1):27.
7. Roberts TJ. 2001. Butterflies of Pakistan. Oxford University Press, London and New York 1-200.
8. Sabir AM, Bhatti AH, Rafi MA, Suhail A. 2000. Distribution of Nymphalid butterflies (brush footed) in districts Rawalpindi and Islamabad. *Pakistan Journal of Biological Sciences*. **3**(8):1253-1254.
9. Munir A, Yasmin N, Rafi MA, Pavulaan H, Wright D. 2007. Bionomic studies of *Papilio demoleus* Linnaeus, the citrus butterfly (Lepidoptera: Papilionidae) from lower Sindh, Pakistan. Taxonomic report of the International Lepidoptera survey **6**(8):1-11.
10. Abbas M, Rafi MA, Inayatullah M, Pavulaan P. 2002. Taxonomy and distribution of butterflies of the Skardu region, Pakistan. Taxonomic Report, the International Lepidoptera Survey (TILS), USA **3**(9):11-19.
11. Haroon SA, Ahmad T. 2013. Diversity of butterfly fauna of Union Council Koaz Bahram Dheri, Pakistan. *J Entomol Zool Stud*. **1**(6):113-117.
12. Rahman UR, Mahmood K, Arif T. 2011. Murree Biodiversity Park-Baseline Report on Entomo fauna. IUCN
13. Pakistan, Islamabad, Pakistan. 2015. Perveen FK, Haroon. Checklist of butterfly (Insecta: Lepidoptera) fauna of Tehsil Tangi, Khyber Pakhtunkhwa, Pakistan. *Arthropods*. **4**(4):98-106.

14. Perveen F, Ahmad A. 2012. Checklist of butterfly fauna of Kohat, Khyber Pakhtunkhwa, Pakistan. *Arthropods*. **1**(3):112-117.
15. De Rhe-Philipe, G.V. 1917. The Butterflies of Lahore. *J. Bombay nat. Hist. Soc.* **25**(1):136-142.
16. Iqbal M, Malik FM, Hussain M, Asharf H, Sarwar KM, Azam I, Umar M. 2016. Butterfly diversity; district Gujrat, Punjab, Pakistan. *Journal of Biodiversity and Environmental Sciences*; **9**(2): 235-243.

**Citation:** Muhammad Amir Sial 2023. Diversity of Butterflies (Insecta: Lepidoptera) in Kasur district, Punjab, Pakistan. *International Journal of Agricultural and Applied Sciences*, 4(2):53-57.

<https://doi.org/10.52804/ijaas2023.427>

**Copyright:** © Muhammad Amir Sial 2023. Creative Commons Attribution 4.0 International License. IJAAS allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.

